Product name: Difenoconazole 250 g/L EC

Product Specification:

| Item | Specification | | |
|---------------------------------------|---------------|--|--|
| Difenoconazole content, g/L | 250 ± 15 | | |
| Water, % | 0.5 max | | |
| pH value | 5.0 – 8.0 | | |
| Emulsion Stability (Dilute 200 times) | Qualified | | |
| Stability at 0 °C | Qualified | | |
| Stability at elevated temperature | Qualified | | |

Introduction:

Difenoconazole is a fungicide, with systemic action. It is a sterol demethylation inhibitor, inhibits the biosynthesis of cell wall sterols, interferes with the normal growth of bacteria, and has a strong inhibitory effect on sporogenesis of plant pathogenic bacteria. This product has a good effect on banana leaf spot.

Direction for use:

| Crop | Target | Application Time | Dosage rate | Method |
|--------|---------------|------------------|------------------|--------|
| Banana | Leaf spot | Before or at the | Dilute 2000-3000 | Spray |
| Banana | Seab | beginning of the | folds | Spray |
| Rice | Sheath blight | disease | 300 – 450 mL/ha | Spray |

Note:

- 1. Do not use in wind or 1 hour before rain.
- 2. Do not mix with alkaline pesticides.
- 3. Fungicides in other mechanism are recommended to use alternately in order to delay resistance.
- 4. Apply three times at most for every banana cycle and the PHI is 42 days. Apply twice at most for every rice cycle and the PHI is 30 days.