

---

**Product name:** Pyraclostrobin 30% SC

**Product Specification:**

Item	Specification
Pyraclostrobin content, %	30 ± 1.5
Suspensibility, %	90 max
pH value	6.0 – 9.0
Pourability, residue after pouring, %	5.0 max
Pourability, residue after washing, %	0.5 max
Persistent foam, mL (after 1min)	30 max
Wet Sieve Test (pass through 75µm sieve), %	98 min
Stability at 0 °C	Qualified
Stability at elevated temperature	Qualified

**Introduction:**

Pyraclostrobin is a fungicide with protectant, curative, and translaminar properties. It can quickly penetrate the leaf, diffuse on the leaf epidermis and adhere closely to the wax layer to form a settling drug film. The effect is stable, rapid and lasts a long time.

**Direction for use:**

Crop	Target	Application Time	Dosage rate	Method
Matrimony vine	Powdery mildew	At the early stage of the disease	Dilution 1500-2500 folds	Spray
Citrus	Anthracoese		Dilution 1500-2000 folds	Spray
Banana	Scab		Dilution 1200-2000 folds	Spray

**Note:**

1. Do not use in wind or 1 hour before rain.
2. Fungicides in other mechanisms are recommended to use by turns to delay resistance.
3. Apply three times at most for every crop cycle. Spray every 15 days. PHI of matrimony vine, citrus and banana is 7 days, 21 days and 42 days respectively.