
Product name: Pyraclostrobin 250g/L EC

Product Specification:

| Item | Specification |
|-----------------------------------|---------------|
| Pyraclostrobin content, g/L | 250 ± 15 |
| Water, % | 0.5 max |
| pH value | 4.0 – 7.0 |
| Persistent foam, mL (after 1min) | 60 max |
| Emulsion stability | Qualified |
| Stability at 0 °C | Qualified |
| Stability at elevated temperature | Qualified |

Introduction:

Pyraclostrobin is a fungicide with protectant, curative, and translaminar properties. It can quickly penetrate the leaf, diffuse on the leaf epidermis and adhere closely to the wax layer to form a settling drug film. The effect is stable, rapid and lasts a long time.

Direction for use:

| Crop | Target | Application Time | Dosage rate | Method |
|------------|--------------|---|-----------------------------|--------|
| Cucumber | Downy mildew | At the early stage of the disease | 600 – 900 mL/ha | Spray |
| Watermelon | Anthracnose | | 300 – 450 mL/ha | Spray |
| Banana | Leaf spot | Before or at the early stage of the disease | Dilution 1000-2000 folds | Spray |

Note:

1. Do not use in wind or 1 hour before rain.
2. Spray evenly and thoughtfully to ensure the prevention effect.
3. Fungicides in other mechanisms are recommended to use by turns to delay resistance.
4. Apply three times at most for every crop cycle. Spray every 10 days.