Product name: Azoxystrobin 200g/L + Thifluzamide 100g/L SC

Product Specification:

Item	Specification	
Azoxystrobin content, g/L	200 ± 12	
Thifluzamide content, g/L	100 ± 10	
Suspensibility of azoxystrobin, %	90 min	
Suspensibility of thifluzamide, %	90 min	
pH value	5.0 – 9.0	
Wet sieve test, % (pass through 75µm sieve)	98 min	
Pourability, residue after pouring, %	5.0 max	
Pourability, residue after washing, %	0.5 max	
Persistent foam, mL (after 1min)	60 max	
Stability at 0 °C	Qualified	
Stability at elevated temperature	Qualified	

Introduction:

Azoxystrobin is a fungicide with protectant, curative, eradicant, translaminar and systemic properties. It inhibits the respiration of the pathogen's mitochondria to prevent its energy synthesis. Thiffuzamide is a succinate dehydrogenase inhibitor in the tricarboxylic acid cycle. It can be rapidly absorbed by roots and leaves and translocated throughout the plant, when applied.

Direction for use:

Crop Target	Application Time	Dosage rate	Method
Rice Sheath blight disease	Before or at the early stage of the disease	450-600 mL/ha	Spray

Note:

- 1. Do not mix with alkaline pesticides.
- 2. Fungicides in other mechanism are recommended to use alternately in order to delay resistance.
- 3. Apply twice at most for every crop cycle. PHI is 21 days for rice.