
Product name: Bispyribac-sodium 400g/L SC

Product Specification:

Item	Specification
Bispyribac-sodium content, g/L	400 ± 20
Suspensibility, %	90 min
pH value	6.5 – 9.5
Wet sieve test, % (pass through 75µm sieve)	98 min
Pourability, residue after pouring, %	5.0 max
Pourability, residue after washing, %	0.5 max
Persistent foam, mL (after 1min)	30 max
Stability at 0 °C	Qualified
Stability at elevated temperature	Qualified

Introduction:

This product is a selective, systemic post-emergence herbicide, absorbed by foliage and roots. It has good control of grasses, sedges and broad-leaved weeds in direct-seeded rice.

Direction for use:

Crop	Target	Application Time	Dosage rate	Method
Direct-seeded rice	Annual weeds and some perennial weeds	At the rice 4 leaf stage or weeds 2-5 leaf stage	75 – 105 mL/ha	Spray

Note:

1. It is forbidden to be used in glutinous rice fields. It is not recommended to use in waseda rice field, because the temperature is low when early rice is used, and it is easy to produce drug damage.
2. After applying in the japonica rice field, the leaves turn yellow, but it recovers within 4-5 days, without affecting the rice yield.
3. Drainage before application, make more than 2/3 of weed stems and leaves above the water surface, irrigation within 24 hours to 72 hours after application, keep 3-5cm water layer in the field for 5-7 days. The water layer should not immerse the rice seedling heart.
4. Try to apply when there is no wind or rain to avoid the drift of fog droplets and harm the surrounding crops.
5. Apply once at most for every crop cycle.