
Product name: Cyhalofop-butyl 50g/L +Penoxsulam 10 g/L OD

Product Specification:

Item	Specification
Cyhalofop-butyl content, g/L	50 ± 5
Penoxsulam content, g/L	10 ± 1.5
Suspensibility of cyhalofop-butyl, %	80 min
Suspensibility of penoxsulam, %	80 min
pH value	5.0 – 8.0
Wet sieve test, % (pass through 75µm sieve)	98 min
Pourability, residue after pouring, %	5.0 max
Pourability, residue after washing, %	0.5 max
Persistent foam, mL (after 1min)	25 max
Stability at 0 °C	Qualified
Stability at elevated temperature	Qualified

Introduction:

This product is a mixture of cyhalofop-butyl and penoxsulam, it has a good control effect on chinensis, barnyard grass (including rice barnyard grass), annual broadleaf grass and sedge in paddy fields.

Direction for use:

Crop	Target	Application Time	Dosage rate	Method
Direct seeding paddy field	Annual weeds	Weeds 2-3 leaf stage	1.5-2.4 L/ha	Spray onto the stem and leaf
Transplant paddy field				

Note:

1. Drainage before application, make more than 2/3 of weed stems and leaves above the water surface, irrigation within 24 hours to 72 hours after application, keep 3-5cm water layer in the field for 5-7 days.
2. During application and 1-2 days after application, the water layer should not immerse the rice seedling heart.
3. If the minimum temperature is below 15°C one week before and after application, or the temperature drops significantly above 5°C within 5 days after application, there will be a risk of phytotoxicity and it is not recommended to use.
3. Apply once at most for every crop cycle.