
Product name: Chlorpyrifos 500 g/L + Cypermethrin 50 g/L EC

Product Specification:

Item	Specification
Chlorpyrifos content, g/L	500 ± 25
Cypermethrin content, g/L	50 ± 5
Water, %	0.5 max
pH value	4.0 – 7.0
Persistent foam, mL (after 1min)	20 max
Emulsion stability	Qualified
Stability at 0 °C	Qualified
Stability at elevated temperature	Qualified

Introduction:

This product is an insecticide composed of Chlorpyrifos and Cypermethrin, with stomach, contact and respiratory action. The product has a long duration of efficacy and rapid efficacy, and can effectively control the *Conopomorpha sinensis Bradley*, cotton bollworm, psyllid and *Bactrocera minax*.

Direction for use:

Crop	Target	Application Time	Dosage rate	Method
Citrus trees	Psyllid	At the peak of young nymph	Dilution 1000 -1500 folds	Spray
Citrus trees	<i>Bactrocera minax</i>	At the peak of young larvae	Dilution 800 -1000 folds	Spray
Litchi trees	<i>Conopomorpha sinensis Bradley</i>		Dilution 1000 -1500 folds	Spray
Cotton	Cotton bollworm		750-1125 g/ha	Spray

Note:

1. Do not use it on vegetables.
2. Do not mix with alkaline pesticides.
3. Do not use in wind or 1 hour before rain.
4. This product is sensitive to melon, lettuce seedling and tobacco. Avoid drifting to these crops to avoid phytotoxicity.
5. Insecticides in other mechanisms are recommended to use by turns to delay resistance.
6. Apply once at most for citrus trees and PHI is 28 days. Apply three times at most for litchi trees and PHI is 28 days. Apply three times at most for cotton and PHI is 21 days.