
Product name: Chlorfenapyr 360 g/L SC

Product Specification:

Item	Specification
Chlorfenapyr content, g/L	360 ± 18
Spontaneity of dispersion, %	80 min
Suspensibility, %	70 min
pH value	5.0 – 8.5
Wet sieve test, % (pass through 75µm sieve)	98 min
Pourability, residue after pouring, %	5.0 max
Pourability, residue after washing, %	0.5 max
Persistent foam, mL (after 1min)	50 max
Stability at 0 °C	Qualified
Stability at elevated temperature	Qualified

Introduction:

This product is pyrrole insecticides, with contact and stomach action. It has strong permeability on the leaf surface and has a certain systemic action. This product has a good control effect on cabbage diamondback moth, scallion thrips and asparagus thrips.

Direction for use:

Crop	Target	Application Time	Dosage rate	Method
Scallion	Thrips	At the early vigorous stage	150 – 195 mL/ha	Spray
Cabbage	Diamondback moth	At the peak of egg hatching or early larval stage	210 – 300 mL/ha	Spray
Asparagus	Thrips	At the early vigorous stage	300 – 500 mL/ha	Spray

Note:

1. Do not mix with alkaline pesticides.
2. Do not use in wind or 1 hours before rain.
3. Insecticides in other mechanism are recommended to use alternately in order to delay resistance.
4. Apply once at most for every crop cycle and PHI is 10 days for scallion, 21 days for cabbage, 3 days for asparagus.