
Product name: Cypermethrin 50 g/L EC

Product Specification:

Item	Specification
Cypermethrin content, g/L	50 ± 5
Cis- isomer content, %	40 - 60
Water, g/kg	2 max
Acidity, g/kg (calculated as H ₂ SO ₄)	1.5 max
Emulsion Stability (Dilute 20 times)	Qualified
Stability at 0 °C	Qualified
Stability at elevated temperature	Qualified

Introduction:

This product is a pyrethroid insecticide with contact and stomach action. It has killing effect on some insect eggs and can control the pests that are resistant to organophosphorus pesticides. It is used control the brassicaceous vegetable cabbage caterpillar, cotton bollworm and cotton aphid.

Direction for use:

Crop	Target	Application Time	Dosage rate	Method
Brassicaceous vegetable	Cabbage caterpillar	Young larva stage	600 – 900 mL/ha	Spray
Cotton	Cotton aphid		900 – 1800 mL/ha	Spray
Cotton	Cotton bollworm		900 – 1800 mL/ha	Spray

Note:

1. Do not use in wind or 1 hour before rain.
2. Insecticides in other mechanism are recommended to use alternately in order to delay resistance.
3. Apply three times at most for every crop cycle. The PHI is 5 days for Brassicaceous vegetable and 7 days for cotton.