
Product name: Abamectin 18 g/L EC

Product Specification:

Item	Specification
Abamectin content, g/L	18 ± 3
α(B _{1a} /B _{1b})	10 min
pH value	4.5 – 7.0
Emulsion Stability (Dilute 20 times)	Qualified
Persistent foam, mL (after 1min)	60 max
Stability at 0 °C	Qualified
Stability at elevated temperature	Qualified

Introduction:

This product is an antibiotic insecticide, with contact and stomach action to insects and mites, and a certain fumigation effect. It has a strong osmotic effect on leaves and can kill pests under the epidermis.

Direction for use:

Crop	Target	Application Time	Dosage rate	Method
Citrus tree	Red spider mite	Egg hatching peak to young larva stage	Dilute 2000-4000 folds	Spray
Pear	Psyllid		Dilute 1500-3000 folds	Spray
Cabbage	Diamond back moth		450 – 600 mL/ha	Spray
Rice	Rice leaf folder		450 – 600 mL/ha	Spray

Note:

1. Do not mix with alkaline pesticides. Insecticides in other mechanism are recommended to use alternately in order to delay resistance.
2. Do not use in wind or 1 hour before rain.
3. Apply twice at most for every crop cycle. The PHI is 14 days for citrus tree and pear, 3 days for cabbage, and 21 days for rice.