

---

**Product name:** Emamectin Benzoate 5% WG

**Product Specification:**

Item	Specification
Emamectin Benzoate content, %	5 ± 0.5
Water, %	3.0 max
Wetting Time, s	90 max
pH value	5.0 – 8.0
Wet sieve test, % (pass through 75µm sieve)	98 min
Suspensibility, %	70 min
Persistent foam, mL (after 1min)	60 max
Attrition resistance, %	90 min
Stability at elevated temperature	Qualified

**Introduction:**

This product is insecticide with contact and stomach action. This product has no internal absorption, but can effectively penetrate into the epidermis tissue of crops, with a long duration of efficacy.

**Direction for use:**

Crop	Target	Application Time	Dosage rate	Method
Cabbage	Beet armyworm	At the peak of young larva	45-75 g/ha	Spray
Rice	Rice leaf folder	At the peak of egg hatching to young larva	150-225 g/ha	Spray
Ginger	Corn borer	From spawning to early hatching	90-150 g/ha	Spray

**Note:**

1. Do not use in wind or 1 hour before rain.
2. Do not mix with alkaline pesticides.
3. Insecticide in other mechanisms are recommended to use by turns to delay resistance.
4. Apply twice at most for cabbage and PHI is 7 days. Apply three times at most for rice and PHI is 14 days. Apply once at most for ginger and PHI is 14 days.