

---

**Product name:** Emamectin Benzoate 10% + Lufenuron 40% WG

**Product Specification:**

Item	Specification
Emamectin Benzoate content, %	10 ± 1.0
Lufenuron content, %	40 ± 2.0
Water, %	2.0 max
pH value	7.0 – 10.0
Wetting Time, s	60 max
Dispersibility, %	80 min
Suspensibility of Emamectin Benzoate, %	80 min
Suspensibility of Lufenuron, %	80 min
Persistent foam, mL (after 1min)	60 max
Attrition resistance, %	95 min
Dustiness, mg	30 max
Stability at elevated temperature	Qualified

**Introduction:**

This product is an insecticide composed of Emamectin benzoate and Lufenuron, with stomach and contact action. It achieves insecticidal effect by affecting the formation of insect epidermis, and can also hinder the transmission of nerve information of pests and kill pests. This product has a good control effect on cabbage caterpillars.

**Direction for use:**

Crop	Target	Application Time	Dosage rate	Method
Cabbage	Cabbage caterpillar	At the peak of young larvae	67.5-135 g/ha	Spray

**Note:**

1. Do not use in wind or 1 hour before rain.
2. Insecticides in other mechanisms are recommended to use by turns to delay resistance.
3. Apply twice at most for every crop cycle. PHI is 7 days.