

---

**Product name:** Thiocyclam-hydrogenoxalate 50% SP

**Product Specification:**

Item	Specification
Thiocyclam-hydrogenoxalate content, %	50 ± 2.5
Water, %	2.0 max
pH value	2.0-5.0
Wetting Time, s	15 max
Persistent foam, mL (after 1min)	50 max
Degree of dissolution and solution stability, % (after 5 min)	2.5 max
Degree of dissolution and solution stability, % (after 18 h)	1.5 max
Stability at elevated temperature	Qualified

**Introduction:**

This product is a neretotoxin insecticide, with contact, stomach, inhalation and fumigation action, and can kill eggs. It has good control effect on rice pests.

**Direction for use:**

Crop	Target	Application Time	Dosage rate	Method
Rice	Rice leaf folder	At the peak of egg hatching to young larva stage	750-1500 g/ha	Spray
Rice	Striped stem borer			Spray
Rice	Yellow stem borer			Spray

**Note:**

1. Beans, cotton, fruit trees are sensitive to thiocyclam-hydrogenoxalate and should not be used.
2. Do not use in wind or 1 hour before rain.
3. Insecticide in other mechanisms are recommended to use by turns to delay resistance.
4. Apply three times at most for every cycle of rice, PHI of rice is 15 days.