
Product name: Indoxacarb 80g/L + Emamectin Benzoate 20g/L SC

Product Specification:

Item	Specification
Indoxacarb content, g/L	80 ± 8
Emamectin Benzoate content, g/L	20 ± 3
Suspensibility of indoxacarb, %	80 min
Suspensibility of emamectin benzoate, %	80 min
pH value	6.0 – 9.0
Wet sieve test, % (pass through 75µm sieve)	98 min
Pourability, residue after pouring, %	5.0 max
Pourability, residue after washing, %	0.5 max
Persistent foam, mL (after 1min)	60 max
Stability at 0 °C	Qualified
Stability at elevated temperature	Qualified

Introduction:

This product is a low toxicity insecticide composed of biogenic pesticide Emamectin Benzoate and chemical pesticide indoxacarb. It has a strong contact and stomach action on pests and can quickly block the transmission of insect nerve impulse, resulting in the target insect movement disorders, paralysis, refusal to feed, and finally death.

Direction for use:

Crop	Target	Application Time	Dosage rate	Method
Rice	Rice leaf folder	Hatching vigorous stage to young larva stage	300-450 mL/ha	Spray
Cabbage	Beet armyworm	At the peak of young larvae	300-450 mL/ha	Spray

Note:

1. Do not mix with alkaline pesticides.
2. Do not use in wind or 1 hour before rain.
3. Insecticides in other mechanisms are recommended to use by turns to delay resistance.
4. Apply twice at most for rice and PHI is 21 days. Apply twice at most for cabbage and PHI is 7 days.